

The newest methods of celestial mechanics

Alexander D. Bruno
abruno@keldysh.ru

Keldysh Institute of Applied Mathematics of RAS

PCA-2020, October 12–17, 2020

Abstract

The last volume of the book "Les méthodes nouvelles de la Mécanique céleste" by [Poincaré, 1899] was published more than 120 years ago. Since then, the following methods have arisen.

- 1 Method of normal forms, allowing to study regular perturbations near a stationary solution, near a periodic solution, near an invariant torus and so on.
- 2 Method of truncated systems, found with a help of the Newton polyhedrons, allowing to study singular perturbations.
- 3 Method of generating families of periodic solutions and invariant tori (regular and singular).
- 4 Method of generalized problems, allowing bodies with negative masses.
- 5 Computation of a net of families of periodic solutions and of invariant tori as a "skeleton" of a part of the phase space.

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method

3.2 The Hill problem

4. Generalized problems

5. Sceletons

6. References

1. Normal forms (1)

Let us consider the Hamiltonian system

$$\dot{\xi}_j = \frac{\partial \gamma}{\partial \eta_j}, \quad \dot{\eta}_j = -\frac{\partial \gamma}{\partial \xi_j}, \quad j = 1, \dots, n \quad (1)$$

with n degrees of freedom in a vicinity of the stationary solution

$$\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) = 0, \quad \boldsymbol{\eta} = (\eta_1, \dots, \eta_n) = 0. \quad (2)$$

If the Hamiltonian function $\gamma(\boldsymbol{\xi}, \boldsymbol{\eta})$ is analytic in the point (2), then it is expanded into the power series

$$\gamma(\boldsymbol{\xi}, \boldsymbol{\eta}) = \sum \gamma_{\mathbf{p}\mathbf{q}} \boldsymbol{\xi}^{\mathbf{p}} \boldsymbol{\eta}^{\mathbf{q}}, \quad (3)$$

where $\mathbf{p} = (p_1, \dots, p_n)$, $\mathbf{q} = (q_1, \dots, q_n) \in \mathbb{Z}^n$, $\mathbf{p}, \mathbf{q} \geq 0$, $\boldsymbol{\xi}^{\mathbf{p}} = \xi_1^{p_1} \xi_2^{p_2} \dots \xi_n^{p_n}$. Here $\gamma_{\mathbf{p}\mathbf{q}}$ are constant coefficients.

1. Normal forms (2)

As the point (2) is stationary, then the expansion (3) begins from quadratic terms. They correspond to the linear part of the system (1). Eigenvalues of its matrix are decomposed in pairs:

$$\lambda_{j+n} = -\lambda_j, \quad j = 1, \dots, n.$$

Let $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_n)$. The canonical changes of coordinates

$$(\boldsymbol{\xi}, \boldsymbol{\eta}) \longrightarrow (\mathbf{x}, \mathbf{y}) \quad (4)$$

preserve the Hamiltonian structure of the system. Here $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_n)$.

1. Normal forms (3)

Theorem ([Bruno, 1972, §12])

There exists a formal canonical invertible transformation (4), bringing the system (1) to the normal form

$$\dot{x}_j = \frac{\partial g}{\partial y_j}, \quad \dot{y}_j = -\frac{\partial g}{\partial x_j}, \quad j = 1, \dots, n, \quad (5)$$

where the series

$$g(\mathbf{x}, \mathbf{y}) = \sum g_{\mathbf{p}\mathbf{q}} \mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}} \quad (6)$$

contains only resonance terms with

$$\langle \mathbf{p} - \mathbf{q}, \boldsymbol{\lambda} \rangle = 0,$$

and the square part $g_2(\mathbf{x}, \mathbf{y})$ has its own normal form (i.e. the matrix of the system is the Hamiltonian analog of the Jordan normal form).

1. Normal forms (4)

If $\lambda \neq 0$, then the normal form (5) is equivalent to a system with smaller number of degrees of freedom and with additional parameters. The normalizing transformation (4) conserves small parameters and linear automorphisms of the initial system (1)

$$(\xi, \eta) \longrightarrow (\tilde{\xi}, \tilde{\eta}), \quad t \rightarrow \tilde{t}.$$

Local families of periodic solutions satisfy the system of equations

$$\frac{\partial g}{\partial y_j} = \lambda_j x_j a, \quad \frac{\partial g}{\partial x_j} = \lambda_j y_j a, \quad j = 1, \dots, n,$$

where a is a free parameter.

1. Normal forms (5)

For the real initial system (1), the coefficients g_{pq} of the complex normal form (5), (6) satisfy to special properties of reality and after a standard canonical linear change of coordinates $(x, y) \rightarrow (X, Y)$ the system (5) transforms in a real system [Bruno, 1994, Ch. I].

There are several methods of computation of coefficients g_{pq} of the normal form (5), (6). The most simple method was described in the book [Zhuravlev (et al.), 2015].

Normal forms of periodic Hamiltonian systems was described in the papers [Bruno, 2020b,c], see also [Bruno, 1994, Ch. II]. Normal forms near a periodic solution, near an invariant torus and near family of them see in [Bruno, 1994, Chs. II, VII, VIII], [Bruno, 1989, Part II].

1. Normal forms (6)

It was shown that each periodic solution belongs to a family of them, but only strongly unstable invariant torus belongs to a family of invariant tori, including torus of maximal dimension [Bruno, 2020a]

Normal form is useful in study stability, bifurcations and asymptotic behavior of solutions.

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method

3.2 The Hill problem

4. Generalized problems

5. Sceletons

6. References

2.1. Truncated Hamiltonian function (1)

Let $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_n)$ and $\boldsymbol{\mu} = (\mu_1, \dots, \mu_s)$ be canonical variables and small parameters respectively. Let a Hamiltonian function be

$$h(\mathbf{x}, \mathbf{y}, \boldsymbol{\mu}) = \sum h_{\mathbf{pqr}} \mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}} \boldsymbol{\mu}^{\mathbf{r}} \quad (7)$$

where $\mathbf{p} = (p_1, \dots, p_n)$, $\mathbf{x}^{\mathbf{p}} = x_1^{p_1} \dots x_n^{p_n}$ and $h_{\mathbf{pqr}}$ are constant coefficients.

2.1. Truncated Hamiltonian function (2)

To each term of sum (7) we put in correspondence its vectorial power exponent $Q = (\mathbf{p}, \mathbf{q}, \mathbf{r}) \in \mathbb{R}^{2n+s}$. Set \mathbf{S} of all points Q with $h_Q \neq 0$ in sum (7) is called as **support** $\mathbf{S} = \mathbf{S}(f)$ of the sum (7). The convex hull $\Gamma(\mathbf{S}) = \Gamma(f)$ of the support \mathbf{S} is called as the **Newton polyhedron** of the sum (7). Its boundary consists of vertices $\Gamma_j^{(0)}$, edges $\Gamma_j^{(1)}$ and faces $\Gamma_j^{(d)}$ of dimensions d : $1 < d \leq 2n + s - 1$. Intersection $\mathbf{S} \cap \Gamma_j^{(d)} = \mathbf{S}_j^{(d)}$ is the boundary subset of set \mathbf{S} .

2.1. Truncated Hamiltonian function (3)

To each *generalized face* $\Gamma_j^{(d)}$ (including vertices and edges) there correspond:

- *normal cone* $\mathbf{U}_j^{(d)}$ in space \mathbb{R}_*^{2n+s} , which is dual to space \mathbb{R}^{2n+s} ;
- *truncated sum*

$$\hat{h}_j^{(d)} = \sum h_{\mathbf{pqr}} \mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}} \boldsymbol{\mu}^{\mathbf{r}} \text{ over } Q = (\mathbf{p}, \mathbf{q}, \mathbf{r}) \in \mathbf{S}_j^{(d)}.$$

It is the first approximation to the sum (7), when $(\log |x_1|, \dots, \log |x_n|, \log |y_1|, \dots, \log |y_n|, \log |\mu_1|, \dots, \log |\mu_s|) \rightarrow \infty$ near $\mathbf{U}_j^{(d)}$.

So by truncated Hamiltonian function we can describe the approximate problems.

2.2 Restricted 3-body problem (1)

Let the two bodies \mathbf{P}_1 and \mathbf{P}_2 with masses $1 - \mu$ and μ respectively turn in circular orbits around their common mass center with the period T . The plane circular restricted three-body problem consists in the study of the plane motion of the body \mathbf{P}_3 of infinitesimal mass under the influence of the Newton gravitation of bodies \mathbf{P}_1 and \mathbf{P}_2 . In the rotating (synodical) standardized coordinate system the problem is described by the Hamiltonian system with two degrees of freedom and with one parameter μ .

2.2 Restricted 3-body problem (2)

The Hamiltonian function has the form

$$h \stackrel{\text{def}}{=} \frac{1}{2} (y_1^2 + y_2^2) + x_2 y_1 - x_1 y_2 - \frac{1 - \mu}{\sqrt{x_1^2 + x_2^2}} - \frac{\mu}{\sqrt{(x_1 - 1)^2 + x_2^2}} + \mu x_1. \quad (8)$$

Here the body $\mathbf{P}_1 = \{X, Y : x_1 = x_2 = 0\}$

and the body $\mathbf{P}_2 = \{X, Y : x_1 = 1, x_2 = 0\}$,

where $X = (x_1, x_2)$, $Y = (y_1, y_2)$. We consider the small values of the mass ratio $\mu \geq 0$.

When $\mu = 0$ the problem turns into the two-body problem for \mathbf{P}_1 and \mathbf{P}_3 . But here the points corresponding to collisions of the bodies \mathbf{P}_2 and \mathbf{P}_3 must be excluded from the phase space. The points of collisions split in parts solutions to the two-body problem for \mathbf{P}_1 and \mathbf{P}_3 .

2.2 Restricted 3-body problem (3)

For small $\mu > 0$ there is a singular perturbation of the case $\mu = 0$ near the body \mathbf{P}_2 . In order to find all the first approximations to the restricted three-body problem, it is necessary to introduce the local coordinates near the body \mathbf{P}_2

$$\xi_1 = x_1 - 1, \xi_2 = x_2, \eta_1 = y_1, \eta_2 = y_2 - 1$$

and to expand the Hamiltonian function in these coordinates.

2.2 Restricted 3-body problem (4)

After the expansion of $1/\sqrt{(\xi_1 + 1)^2 + \xi_2^2}$ in the Maclaurin series, the Hamiltonian function (8) takes the form

$$h + \frac{3}{2} - 2\mu \stackrel{\text{def}}{=} \frac{1}{2}(\eta_1^2 + \eta_2^2) + \xi_2\eta_1 - \xi_1\eta_2 - \xi_1^2 + \frac{1}{2}\xi_2^2 + \\ + f(\xi_1, \xi_2^2) + \mu \left\{ \xi_1^2 - \frac{1}{2}\xi_2^2 - \frac{1}{\sqrt{\xi_1^2 + \xi_2^2}} - f(\xi_1, \xi_2^2) \right\}, \quad (9)$$

where f is the convergent power series, where the terms of order less than three are absent.

2.2 Restricted 3-body problem (5)

Let for each term of sum (9) we put

$$p = \text{ord } \xi_1 + \text{ord } \xi_2, \quad q = \text{ord } \eta_1 + \text{ord } \eta_2, \quad r = \text{ord } \mu.$$

Then support \mathbf{S} of the expansion (9) consists of the points

$$(0, 2, 0), (1, 1, 0), (2, 0, 0), (k, 0, 0), (2, 0, 1), (-1, 0, 1), (k, 0, 1),$$

where $k = 3, 4, 5, \dots$. The convex hull of the set \mathbf{S} is the polyhedron $\Gamma \subset \mathbb{R}^3$. The surface $\partial\Gamma$ of the polyhedron Γ consists of faces $\Gamma_j^{(2)}$, edges $\Gamma_j^{(1)}$ and vertices $\Gamma_j^{(0)}$. To each of the elements $\Gamma_j^{(d)}$ there corresponds the truncated Hamiltonian $\hat{h}_j^{(d)}$, that is the sum of those terms of Series (9), the points $Q = (p, q, r)$ of which belong to $\Gamma_j^{(d)}$.

2.2 Restricted 3-body problem (6)

Fig. 1 shows the polyhedron Γ , which is the semi-infinite trihedral prism with an oblique base. It has four faces and six edges. Let us consider them.





Figure 1 The polyhedron Γ for the Hamiltonian function (9) in coordinates p, q, r .

2.2 Restricted 3-body problem (7)

The face $\Gamma_1^{(2)}$, which is the oblique base of the prism Γ , contains vertices $(0, 2, 0)$, $(2, 0, 0)$, $(-1, 0, 1)$ and the point $(1, 1, 0) \in \mathbf{S}$.



2.2 Restricted 3-body problem (8)

To the face there corresponds the truncated Hamiltonian function

$$\hat{h}_1^{(2)} = \frac{1}{2} (\eta_1^2 + \eta_2^2) + \xi_2 \eta_1 - \xi_1 \eta_2 - \xi_1^2 + \frac{1}{2} \xi_2^2 - \frac{\mu}{\sqrt{\xi_1^2 + \xi_2^2}}. \quad (10)$$

It describes the Hill problem [Hill, 1878], which is a non-integrable one.

The power transformation


$$\tilde{\xi}_i = \xi_i \mu^{-1/3}, \quad \tilde{\eta}_i = \eta_i \mu^{-1/3}, \quad i = 1, 2, \quad (11)$$

reduces the corresponding Hamiltonian system to the Hamiltonian system with the Hamiltonian function of the form (10), where ξ_i, η_i, μ must be substituted by $\tilde{\xi}_i, \tilde{\eta}_i, 1$ respectively.

2.2 Restricted 3-body problem (9)

The face $\Gamma_2^{(2)}$ contains points $(0, 2, 0)$, $(1, 1, 0)$, $(2, 0, 0)$ and $(k, 0, 0) \subset \mathbf{S}$.

To the face there corresponds the truncated Hamiltonian function $\hat{h}_2^{(2)}$, which is obtained from the function h when $\mu = 0$. It describes the two-body problem for \mathbf{P}_1 and \mathbf{P}_3 , which is an integrable one.




2.2 Restricted 3-body problem (10)

The edge $\Gamma_1^{(1)}$. It includes points $(0, 2, 0)$ and $(-1, 0, 1) \in \mathbf{S}$. The corresponding truncated Hamiltonian function is


$$\hat{h}_1^{(1)} = \frac{1}{2}(\eta_1^2 + \eta_2^2) - \frac{\mu}{\sqrt{\xi_1^2 + \xi_2^2}}. \quad (12)$$

It describes the *two-body problem* for \mathbf{P}_2 and \mathbf{P}_3 . The power transformation (11) transforms it into the Hamiltonian system with the Hamiltonian function of the form (12), where ξ_i, η_i, μ must be substituted by $\tilde{\xi}_i, \tilde{\eta}_i, 1$ respectively.



2.2 Restricted 3-body problem (11)

The edge $\Gamma_2^{(1)}$ includes points $(2, 2, 0), (1, 1, 0), (0, 2, 0) \in \mathbf{S}$. To it there corresponds the truncated Hamiltonian function (10) with $\mu = 0$. It describes the intermediate problem (between the Hill problem and the two-body problem for \mathbf{P}_1 and \mathbf{P}_3), which is an integrable one. This first approximation was introduced by [Hénon, 1969].



2.2 Restricted 3-body problem (12)

Thus, the first approximation to the original restricted problem with the Hamiltonian function (9) depends on the distance from the body \mathbf{P}_2 in the following manner:

- very close to the body \mathbf{P}_2 , it is the two-body problem for bodies \mathbf{P}_2 and \mathbf{P}_3 with the Hamiltonian function (12);
- simply close to the body \mathbf{P}_2 , it is the Hill problem with Hamiltonian (10);
- farther from the body \mathbf{P}_2 , it is the intermediate Hénon problem with Hamiltonian (10) and $\mu = 0$;
- and far from the body \mathbf{P}_2 , it is the two-body problem for bodies \mathbf{P}_1 and \mathbf{P}_3 .

2.2 Restricted 3-body problem (13)

Near the body \mathbf{P}_2 , the periodic solutions to the restricted problem are either perturbations of periodic solutions to all four mentioned first approximations or they are results of the matching of the hyperbolic orbits of the two-body problem for \mathbf{P}_2 and \mathbf{P}_3 with arc-solutions to the two-body problem for \mathbf{P}_1 and \mathbf{P}_3 , or to the intermediate problem. In [Benest, 1976; Kogan, 1989; Lidov (et al.), 1990; 1993; 1994] the periodic solutions to the intermediate problem were used as the generating ones in order to find quasi-satellite orbits of the restricted problem.

2.3 Truncated systems (1)

Now we consider the aggregate of polynomials

$$f_1(X), \dots, f_m(X), \quad X \in \mathbb{R}^{m'} \text{ or } \mathbb{C}^{m'}. \quad (13)$$

2.3 Truncated systems (2)

To each f_i there corresponds its support and all the accompanying objects: polyhedrons Γ_j , faces $\Gamma_{jk_j}^{(d_j)}$, normal cones $U_{jk_j}^{(d_j)}$, boundary subsets $S_{jk_j}^{(d_j)}$, truncated polynomials $\hat{f}_{jk_j}^{(d_j)}$. Besides, to each non-empty intersection

$$U_{1k_1}^{(d_1)} \cap \cdots \cap U_{mk_m}^{(d_m)} \quad (14)$$

there corresponds the aggregate of truncations of the form

$$\hat{f}_{1k_1}^{(d_1)}, \dots, \hat{f}_{mk_m}^{(d_m)},$$

which is the first approximation to the aggregate (13), when $\log |X| \rightarrow \infty$ near the intersection (14); and it is named the *truncation of the aggregate* (13).

2.3 Truncated systems (3)

We consider now the system of equations

$$f_j = 0, \quad j = 1, \dots, m, \quad (15)$$

corresponding to the aggregate (13). To System (15) there correspond all objects indicated for the aggregate (13), and also the *truncated systems of equations*

$$\hat{f}_{jk_j}^{(d_j)} = 0, \quad j = 1, \dots, m, \quad (16)$$

each of which corresponds to one aggregate of truncations (6). We say that the truncated system (16) is the *truncation of System (15) with respect to the order* $P \neq 0$ if the vector P lies in the cone (14). Every truncated system (16) is the first approximation to one complete system (15).

2.4 Periodic solutions to periodic Hamiltonian system (1)

Normal form of a periodic Hamiltonian function with n degrees of freedom near zero solution is reduced to a stationary Hamiltonian function

$$h(\mathbf{u}, \mathbf{v}, \mu) = \sum h_{\mathbf{p}\mathbf{q}\mathbf{r}m} \mathbf{u}^{\mathbf{p}} \mathbf{v}^{\mathbf{q}} \mu^{\mathbf{r}}, \quad (17)$$

where $\mathbf{p}, \mathbf{q} \in \mathbb{Z}^n$, $\mathbf{r} \in \mathbb{Z}^s$, $m \in \mathbb{Z}$, $\mathbf{p}, \mathbf{q}, \mathbf{r} \geq 0$ and

$$\langle \boldsymbol{\lambda}, \mathbf{p} - \mathbf{q} \rangle = -im.$$

2.4 Periodic solutions to periodic Hamiltonian system (2)

For $\mu = 0$, expansion of h (17) begins from terms of order 3. Local families of periodic solutions to the initial system correspond to local families of stationary points of the reduced normal form with Hamiltonian (9). These stationary points satisfy system of equations

$$\frac{\partial h}{\partial v_j} = 0, \quad \frac{\partial h}{\partial u_j} = 0, \quad j = 1, \dots, n. \quad (18)$$

To solve the system, we must to consider truncated systems and find their solutions, which gives the first approximations to solutions of the system (18).

Other applications: the Beletskii equation for oscillation of a satellite [Bruno, Varin, 1997]; the problem of periodic orbits with close approach to planet and to Earth [Bruno, 1981].

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method

3.2 The Hill problem

4. Generalized problems

5. Sceletons

6. References

3.1 Method

Let a Hamiltonian function $H(\boldsymbol{\mu})$ analytically depend from small parameters $\boldsymbol{\mu} = (\mu_1, \dots, \mu_s)$ and corresponding Hamiltonian system has sets $\mathcal{M}_j(\boldsymbol{\mu})$ of solutions. Some of these sets can have limits $\mathcal{M}_j(0)$, when $\boldsymbol{\mu} \rightarrow 0$. Sets $\mathcal{F}_j(0)$ are called as **generating**. Their solutions are compositions of parts of solutions of the limit Hamiltonian system with $\boldsymbol{\mu} = 0$. Especially interesting are generating families of periodic solutions and of invariant tori.

If that limit system is integrable, than generating families can be described analytically. That approach was proposed by [Hénon, 1969] for generating families of periodic solutions and was used for the Hill problem, for the restricted three-body problem [Bruno, 1994; Hénon, 1968; Hénon, 1997; 2001], for the Belletskii equation [Bruno, Varin, 1997; 2007].

3.2 The Hill problem (1)

Its Hamiltonian function is

$$H = \frac{1}{2} (\eta_1^2 + \eta_2^2) + \xi_2 \eta_1 - \xi_1 \eta_2 - \xi_1^2 + \frac{1}{2} \xi_2^2 - \frac{1}{\sqrt{\xi_1^2 + \xi_2^2}}. \quad (19)$$

The corresponding system

$$\dot{\xi}_j = \frac{\partial H}{\partial \eta_j}, \quad \dot{\eta}_j = -\frac{\partial H}{\partial \xi_j}, \quad j = 1, 2$$

describes the motion of Moon (\mathbf{P}_3) with zero mass under attraction of Sun (\mathbf{P}_1) disposed at infinity and Earth (\mathbf{P}_2) with mass 1 disposed in origin. Hamiltonian (19) is analytic in

$$\boldsymbol{\xi}, \boldsymbol{\eta} \in \mathbb{R}^4 \setminus \{\xi_1 = \xi_2 = 0\}.$$

3.2 The Hill problem (2)

We make canonical transformation of coordinates

$$\xi_j = \varepsilon X_j, \quad \eta_j = \varepsilon Y_j, \quad j = 1, 2.$$

Then we obtain the Hamiltonian system

$$\dot{X}_j = \frac{\partial h}{\partial Y_j}, \quad \dot{Y}_j = -\frac{\partial h}{\partial X_j}, \quad j = 1, 2, \quad (20)$$

where

$$h = \frac{1}{2} (Y_1^2 + Y_2^2) + X_2 Y_1 - X_1 Y_2 - X_1^2 + \frac{1}{2} X_2^2 - \frac{1}{\varepsilon^3 \sqrt{X_1^2 + X_2^2}}.$$

3.2 The Hill problem (3)

We put $\varepsilon = \sqrt{2|H|}$ and $H \rightarrow -\infty$. Then in limit we obtain system (20) with

$$h = h_0 = \frac{1}{2} (Y_1^2 + Y_2^2) + X_2 Y_1 - X_1 Y_2 - X_1^2 + \frac{1}{2} X_2^2.$$

It is the Hénon's problem. For h_0 system (20) is linear and hence integrable. It is enough to consider it for $h_0 = \frac{1}{2}$. It has one regular periodic solution

$$X_1(t) = \cos t, \quad X_2(t) = -2 \sin t.$$


3.2 The Hill problem (4)

If the orbit $(X_1(t), X_2(t))$ of a solution of the Hénon problem comes through the point


$$X_1 = X_2 = 0, \tag{21}$$

then the body P_3 collides with body P_2 and the solution cannot be continued through that collision. So solutions are divided into independent parts by the point (21). Hénon, 1969 found all arc-solutions, which begin and end by such collisions. They form the countable set of two types. The arc-solutions of the first type were denoted by symbol $\pm j$, $j \in \mathbb{N}$, and are epicycloids. In Figure 2 they are shown for $j = 1, 2, 3$.


3.2 The Hill problem (5)



(a) Arc +1



(b) Arc +2




(c) Arc +3

Figure 2 Arc-solutions of the first type j : +1, +2 and +3.


3.2 The Hill problem (6)

The arcs with negative values of j are symmetric with respect to the axis X_2 . The arc-solutions of the second type are denoted by symbols i and e and their orbits are ellipses passing through the origin (Figure 3).

3.2 The Hill problem (7)



(a) Arc i



(b) Arc e

Figure 3 Arc-solutions of the second type i and e .

3.2 The Hill problem (8)

A sequence of arc-solutions which does not contain two identical arcs of the second type in succession is a generating solution and it is called *generating sequence* [Batkhin, 2013] for the Hill problem. All known families of periodic solutions of the Hill problem include at most one generating sequence.

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method

3.2 The Hill problem

4. Generalized problems

5. Sceletons

6. References

4. Generalized problems

Usually in celestial mechanics we consider bodies with non-negative masses. But [Batkhin, 2014] proposed to consider problems, where some masses are negative. In the Hill problem with mass of the body \mathbf{P}_2 equal to -1 (so-called *anti-Hill problem*), families of periodic solutions are continuations of families of with periodic solutions of the usual Hill's problem. So computation of families of periodic solutions more convenient to make for both Hill's and anti-Hill's problems. Such approach gave new families for the Hill's problem.

Figure 4 shows diagram of connection between families of the Hill's (left part) and the anti-Hill's problems (right part). Central column gives generating sequences of the families.




Figure 4 Diagram of connection between families.

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method

3.2 The Hill problem

4. Generalized problems

5. Sceletons

6. References

5. Sceletons (1)

In some parts of the phase space of a Hamiltonian system there are a lot of families of periodic solutions and of invariant tori. These families form a “skeleton” of the phase space. So computation of such families is very useful for study the structure of the phase space. [Batkhin, 2019] mentioned that in systems with a finite group of symmetries, the majority of such families consist of periodic solutions, with are invariant under all symmetries of the group.

There are a lot of computed families of periodic solutions in different problems of celestial mechanics, but their number is not enough to form a skeleton. Recent results in that directions for the restricted three-body problem see in [Bruno, Varin, 2007; 2009a,b,c,d; 2011; 2012].

5. Sceletons (2)

A method for computation of a family of invariant tori was proposed by Simo [Simó, 1990], but up today there are a few cases of their computations.

1. Normal forms

2. Newton polyhedrons

2.1 Truncated Hamiltonian function [Bruno2000]

2.2 Restricted 3-body problem [Bruno2000]

2.3 Truncated systems

2.4 Periodic solutions to periodic Hamiltonian system [Bruno, 2020b,c]

3. Generating families of periodic solutions and of invariant tori

3.1 Method







3.2 The Hill problem

4. Generalized problems





5. Sceletons

6. References






References (1)

-  *Batkhin A. B.* Symmetric Periodic Solutions of the Hill's Problem. I. // Cosmic Research. 2013. Vol. 51, no. 4. P. 275–288. DOI: 10.1134/S0010952513040035.
-  *Batkhin A. B.* Web of Families of Periodic Orbits of the Generalized Hill Problem. // Doklady Mathematics. 2014. Vol. 90, no. 2. P. 539–544. DOI: 10.1134/S1064562414060064.
-  *Batkhin A. B.* On the structure of the Hamiltonian phase flow near symmetric periodic solution. // Preprints of KIAM. 2019. No. 69. DOI: 10.20948/prepr-2019-69. (in Russian).
-  *Benest D.* Libration effects for retrograde satellites in the restricted three-body problem. I: Circular plane Hill's case. // Celestial Mechanics. 1976. No. 13. P. 203–215.
-  *Bruno A. D.* Analytical form of differential equations (II). // Trans. Moscow Math. Soc. 1972. Vol. 26. P. 199–239.
-  *Bruno A. D.* On periodic flybys of the Moon. // Celestial Mechanics. 1981. Vol. 24. P. 255–268.






References (2)

-  *Bruno A. D.* Local Methods in Nonlinear Differential Equations. Berlin – Heidelberg – New York – London – Paris – Tokyo: Springer–Verlag, 1989.
-  *Bruno A. D.* The Restricted 3–body Problem: Plane Periodic Orbits. Berlin: Walter de Gruyter, 1994.
-  *Bruno A. D.* Families of periodic solutions and invariant tori of Hamiltonian system without parameters. // Preprints of KIAM. 2020a. No. 71. DOI: 10.20948/prepr-2020-71. URL: http://keldysh.ru/papers/2018/prep2020_71.pdf ; (in Russian).
-  *Bruno A. D.* Normal form of a Hamiltonian system with a periodic perturbation. // Computational Mathematics and Mathematical Physics. 2020b. Vol. 60, no. 1. P. 36–52. DOI: 10.1134/S0965542520010066.






References (3)

-  Bruno A. D. Normalization of the periodic Hamiltonian system. // Programming and Computer Software. 2020c. Vol. 46, no. 2. P. 76–83. DOI: 10.1134/S0361768820020048.
-  Bruno A. D., Varin V. P. The limit problems for the equation of oscillations of a satellite. // Celestial Mechanics and Dynamical Astronomy. 1997. Vol. 67, no. 1. P. 1–40.
-  Bruno A. D., Varin V. P. Periodic solutions of the restricted three-body problem for small mass ratio. // J. Appl. Math. Mech. 2007. Vol. 71, no. 6. P. 933–960.
-  Bruno A. D., Varin V. P. Closed families of periodic solutions of the restricted problem. // Solar System Research. 2009a. Vol. 43, no. 3. P. 253–276.
-  Bruno A. D., Varin V. P. Families c and i of periodic solutions of the restricted problem for $\mu = 5 \cdot 10^{-5}$. // Solar System Research. 2009b. Vol. 43, no. 1. P. 26–40.






References (4)

-  Bruno A. D., Varin V. P. Family h of periodic solutions of the restricted problem for big μ . // Solar System Research. 2009c. Vol. 43, no. 2. P. 158–177.
-  Bruno A. D., Varin V. P. Family h of periodic solutions of the restricted problem for small μ . // Solar System Research. 2009d. Vol. 43, no. 1. P. 2–25.
-  Bruno A. D., Varin V. P. On asteroid distribution. // Solar System Research. 2011. Vol. 45, no. 4. P. 323–329.
-  Bruno A. D., Varin V. P. Periodic solutions of the restricted three body problem for small μ and the motion of small bodies of the Solar system. // Astronomical and Astrophysical Transactions (AApTr). 2012. Vol. 27, no. 3. P. 479–488.
-  Hénon M. Sur les orbites interplanétaires qui rencontrent deux fois la terre. // Bull. astron. Ser. 3. 1968. T. 3, n° 3. P. 377–402.


References (5)

-  *Hénon M.* Numerical exploration of the restricted problem. V. Hill's case: periodic orbits and their stability. // *Astron. & Astrophys.* 1969. Vol. 1. P. 223–238.
-  *Hénon M.* *Generating Families in the Restricted Three-Body Problem.* Berlin, Heidelberg, New York: Springer, 1997. 278 p. (Lecture Note in Physics. Monographs ; 52).
-  *Hénon M.* *Generating Families in the Restricted Three-Body Problem. II. Quantitative Study of Bifurcations.* Berlin, Heidelberg, New York: Springer-Verlag, 2001. (Lecture Note in Physics. Monographs ; 65).
-  *Hill G. W.* *Researches in the Lunar Theory.* // *Amer. J. Math.* 1878. Vol. 1. P. 5–26, 129–147, 245–260.
-  *Kogan A. Y.* Distant satellite orbits in the restricted circular three-body problem. // *Cosmic Research.* 1989. Vol. 26, no. 6. P. 705–710.

References (6)

-  *Lidov M. L., Vashkov'yak M. A.* Analytical Celestial Mechanics. // / ed. by K. V. Kholshchevnikov. Kazan': Kasan' University, 1990. Chap. Quasisatellite periodic orbits. P. 53–57. in Russian.
-  *Lidov M. L., Vashkov'yak M. A.* Perturbation theory and analysis of evolution of quasisatellite orbits in the restricted three-body problem. // Cosmic Research. 1993. Vol. 31, no. 2. P. 187–207.
-  *Lidov M. L., Vashkov'yak M. A.* On quasi-satellite orbits in a restricted elliptic three-body problem. // Astronomy Letters. 1994. Vol. 20. P. 676–690.
-  *Poincaré H.* Les méthodes nouvelles de la mécanique céleste. T. 3. Paris : Gauthier-Villars, 1899.
-  *Simó C.* Modern Methods in Celestial Mechanics. // / ed. by C. F. C. Benest. Editions Frontières, 1990. Chap. Analytical and numerical computation of invariant manifolds. P. 285–330.

References (7)

-  Zhuravlev V. F., Petrov A. G., Shunderyuk M. M. Selected Problems of Hamiltonian Mechanics. Moscow: LENAND, 2015. P. 304. (in Russian).